

THE
AMERICAN JOURNAL
OF THE MEDICAL SCIENCES
FOR JANUARY, 1883.

ARTICLE I.

THE REVIVAL OF SYMPHYSIOTOMY IN ITALY, WITH COMPARATIVE TABLES OF THE EARLY AND LATER CASES, SHOWING THAT THE OPERATION HAS BEEN MORE FREQUENTLY PERFORMED IN THAT COUNTRY IN THE LAST SEVENTEEN YEARS THAN IN ALL EUROPE IN THE PREVIOUS EIGHTY, AND WITH FAR BETTER RESULTS. THE WHOLE SUBJECT EXAMINED HISTORICALLY AND CLINICALLY. By ROBERT P. HARRIS, A.M., M.D., Fellow of the College of Physicians, Phila.; Member of the Obstetrical Society, Phila.; Corresponding Member of the Royal Medico-Chirurgical Academy of Naples, etc.

ONE hundred and fourteen years ago, a French medical student who had but recently entered upon the study of his profession, sent to the Academy of Surgery of Paris, a proposition, in which he advocated a division of the symphysis pubis in labour in cases of deformed pelvis, and claimed that such a section would admit of the enlargement of the pelvic canal, and the delivery of the fœtus. This proposition of Jean René Sigault was ridiculed in the Academy, and treated as the wild scheme of an ignorant youth. Not convinced of his error, Sigault made the plan the subject of a thesis in 1773, and in 1777 made trial of the operation, as shown in table I, case 1. The fœtus being small, and the pelvis of the rachitic dwarf larger than was admitted, there was no strain upon the sacro-iliac synchondroses, and the child was delivered alive. The woman recovered and was exhibited with her infant, to the annoyance of the Academy of Surgery, and the professional advantage of the operator, who was presented with a medal by the Faculty of Medicine, and with a pension by the government; one of the latter being also given to the woman.

Being very unwisely lauded as a public benefactor on the merits of one case, and it neither a crucial one, nor in all respects a perfect success,
No. CLXIX.—JAN. 1883. 2

there was at once created a feeling of opposition to the operator and his scheme, in which the Academy of Surgery was by no means idle, and the medical profession became divided into *Cæsareanists* and *Symphysiotomists*, Sigault having claimed that his operation was to set aside the Cæsarean section in cases of pelvic deformity. The basis of the scheme was the claim, that the pelvic symphyses became relaxed during pregnancy, and could be acted upon like a hinge to a considerable extent without rupture. This was denied, and to prove it an error, the operation was performed upon a number of dead subjects in which, the condition of the body favouring the production of the result anticipated, the ligaments tore in the experiments. Years afterward, when the excitement had passed away, and the operation was rarely performed, it was found that women dying in labour and operated upon soon afterward, exhibited a marked mobility in the sacro-iliac synchondroses, which would admit of a wide separation of the pubic bones without laceration. Dr. Ainsiaux, of Liège, under these circumstances, in 1811, obtained three inches of separation, but found that after the woman had been dead 36, 38, 48, or 54 hours, he could only obtain from $1\frac{1}{4}$ to $1\frac{1}{2}$ inches. Dr. Giraud, of Paris, in 1800, reached the same opinion.

Symphysiotomy, so far as the design and execution of it upon the living woman were concerned, was entirely original with Sigault; but it had been performed post-mortem many years before; first in Warsaw, in 1655, by Jean V. C. Delacourvée, of France, then in practice there, upon a woman of 48, who died after a labour of four days; and second in 1766 by Prof. Jos. Jacques Plenck, of Bude, Hungary, who, finding in a post-mortem Cæsarean section, that the fetal head was locked in the pelvis, opened the symphysis pubis and liberated it. It does not appear to have entered the thoughts of either of these obstetricians that such a section might be applicable to the living parturient woman.

The second Sigaultian operation, as shown by table 1, was performed upon a pubes previously diseased, and therefore unsuitable for the experiment. The fatal termination of the case encouraged the opposition, and brought no credit to the German professor who operated. An examination of the early operations gave very little encouragement to one inclined to hope in the possibility of the scheme supplanting gastro-hysterotomy in deformed pelvis with a very short conjugate diameter, or even being adapted to cases as low as $2\frac{1}{2}$ inches; still Sigault felt inclined to test his plan in an extreme case, as shown in No. 12, and the fatal result was such as might have been anticipated. The best that can be said for him is, that he was in error as to his measurements of the pelvis; otherwise we must regard his act as next to murderous. He was evidently very much disappointed in the capabilities of his operation, and declined to divide the symphysis in his closing days, in a case having a $2\frac{1}{2}$ inch conjugate. In fact, his whole experience amounted to only five cases,

covering a period of one year. He lost one woman, and four children out of five. His associate, Leroy, appeared to have persevered for a longer period, although he had no more operations; he is generally credited with having saved four children; but his correct credit is three. In the year following Sigault's first operation (1778), it will be noticed there were eleven cases, losing six women and ten children. This being quite discouraging, the number in 1779 fell off to four, all in France; and in 1780 to two; these six in succession recovered; still the operation fell in general estimation until the close of the century, when the average became less than one a year. For more than sixty years the operation has been confined exclusively to Italy, and very few operations have been performed outside of the city of Naples. Here it will be noticed there were only twenty-two operations in forty years, 1818-1858, and of these there were fourteen performed by one man, of which nine recovered, with a loss of ten children.

Even in Naples, the mortality of the women and children was such that the operation may be said to have died out in 1858; there were four cases in eighteen years, and then not another until at the end of eight more years the operation was put upon a new trial, by new obstetrical surgeons in the Ospedale dei Incurabili.

With a great deal of care and research, I have been enabled to prepare for the reader and the benefit of future investigators the accompanying historical table of the early symphysiotomies. This in few words will show the character of cases operated upon, and the fatality to women and children. There is no tabular record extant earlier than that of Prof. Alfonso Corradi, of Bologna, in his History "*Dell'Ostetricia in Italia*," which commences with the case of Bruno Amantea in 1807 (No. 35, table 1), and records twenty-four operations in all, which are exclusively Neapolitan. The other forty-six cases are placed in table for the first time, and the chronological order adopted for the whole seventy will greatly facilitate their examination, and comparison with those in table 2, which presents so different a picture that we are inclined at first to doubt the reliability of the statement. This is made, however, by one of eminent position in Naples, as an obstetrician among the wealthy, by which he has made himself independent; a professor in the University of Naples; a hospital surgeon; and a member of the Royal Academy of Medicine and Surgery. Corradi, who is a very critical historian, makes no doubt about the first twenty-six cases in table 2; and my experience with Italian correspondents leads me to place every reliance upon their statements. Besides, what motive could there be in Prof. Morisani to give an inaccurate hospital record as a basis of an elaborate and voluminous pamphlet; of a second brochure sent to the International Medical Congress; and, finally, of a table prepared for me, according to my own plan, from the hospital records and by private research? I make this

defence in advance because there are those who are already inclined to explain away the marvellous improvements in the results of Sigault's operation by doubting their reliability. Prof. Morisani is associated with four of the best known obstetricians and gynecologists in Italy, as a co-editor in an Italian edition of Cazeaux and Tarnier's Midwifery; in the last section of which, just completed, he has committed himself to, and staked his reputation upon, a similar report to that of the operations here presented in my second table. We may not be able fully to comprehend the reason for the great change in the results of the operation of Sigault now, over those which followed the same section, and in the same hospital, under Galbiati; but this is no excuse for doubting the record. The same operators are candid in admitting failures in ovariectomy, the Cesarean section, and Porro's operation; then why would they not be equally so in symphysiotomy? The shoe pinches just here. If Prof. Morisani tells the whole truth, then much that has been said about the dangers and impracticabilities of symphysiotomy in obstetrical text-books will have to be recalled.

General Summary of the 70 Cases in Table 1.—Operations performed in Italy, 31; France, 25; Holland, 4; Belgium, 4; Germany, 3; Spain, 1; England, 1; locality not given, 1 = 70. Women lost, 26; children born dead, or moribund, 47; children saved, 20; fate not ascertained, 3. *Causes of death in the women*, viz., pelvic injuries, 4; the same resulting in gangrene, 6; pelvic cellulitis, 1; putrid fever, 1; anasarca, 1; exhaustion and previous disease, 1; shock, 1; syncope, 1; and not mentioned, 10 = 26. Women delivered of children before the symphysiotomy, or at a later period, 19. Cases operated upon twice each, 4, viz., 4-17, 23-29, 36-38, 37-40. Cases of misdirected incision, cutting the bone instead of the cartilage, 3. Prolapsus uteri is mentioned as having followed the operation in 3 cases; and incontinence of urine from fistula in 1 of these and 3 others. Maternal mortality up to 1858, 37 per cent.; fetal mortality, 67 per cent. Contrast this percentage with that shown in table 2. One-half of the 70 operations were performed by six operators, viz., Prof. Galbiati, 15; Dr. Sigault, 5; Dr. Leroy, 5; Dr. De Cambon, 4; Dr. Gianni, 3; and Dr. Jacolucci, 3 = 35. Of this number, 22 women were saved, and 13 lost; and 24 children perished.

The fetal mortality in the early days of symphysiotomy had much to do with diminishing the frequency of its performance. As the operation was largely advocated in the interest of the fetus, the failure to save it reacted upon the measure as a plan for saving the mother. The question really at issue was between the destruction of the fetus and its delivery, *per vias naturales*, alive. Many of the women had already been delivered of dead children without craniotomy, and some with it, and the object to be gained by separating the symphysis pubis was the saving of the child alive. When this failed in its accomplishment, the crotchet was resorted to, and the fetus destroyed in the interest of the mother.

1. Table of 70 Symphysiotomies, from the first case under Sigault, down to 1858.

No.	Date.	Authority.	Locality.	Conjugate diameter.	Result to woman.	Result to child.	Remarks.
1	Oct. 1, 1777	Dr. Jean René Sigault	Paris	3 in. 6 lines	Recovered	Alive	3 ft 5 in. high; wife of a soldier; 3th pregnancy; other children dead at presentation. Biparietal diam. 3 in. 4 l. Inconspicuous scar on face; no pain; no fever; no inflammation.
2	Feb. 4, 1778	Prof. Siebold	Würzburg	3 in. 9 l.	Died	Lost	Wife of a soldier; had had 7 children, all of whom died; one by embryotomy; pubes ossified, and was saved in labour. Fetus turned for delivery. Woman died of puerperal inflammation.
3	Feb. 21, 1778	Dr. Despres de Monmou	St. Paul de Leon, France		Recovered	Lost	Wife of a soldier; had had 7 children, all of whom died; one by embryotomy; pubes ossified, and was saved in labour. Fetus turned for delivery. Woman died of puerperal inflammation.
4	Mar. 23, 1778	Dr. De Cambon	Mons, Belgium		Recovered	Lost	Wife of a soldier; had had 7 children, all of whom died; one by embryotomy; pubes ossified, and was saved in labour. Fetus turned for delivery. Woman died of puerperal inflammation.
5	Apr. 6, 1778	Dr. A. Kessel de Vauzemo	Paris		Died	Alive	Wife of a soldier; had had 7 children, all of whom died; one by embryotomy; pubes ossified, and was saved in labour. Fetus turned for delivery. Woman died of puerperal inflammation.
6	Apr. 6, 1778	Dr. François J. Nagel	Spire, Bavaria	3 in.	Died	Lost	Wife of a soldier; had had 7 children, all of whom died; one by embryotomy; pubes ossified, and was saved in labour. Fetus turned for delivery. Woman died of puerperal inflammation.
7	Apr. 24, 1778	Dr. L'Escaudé and Rotz	Arras, France	3 in. 0 l.	Died	Lost	Wife of a soldier; had had 7 children, all of whom died; one by embryotomy; pubes ossified, and was saved in labour. Fetus turned for delivery. Woman died of puerperal inflammation.
8	May 10, 1778	Prof. Guérard	Dusseldorf	2 in. 0 l.	Died	Lost	Wife of a soldier; had had 7 children, all of whom died; one by embryotomy; pubes ossified, and was saved in labour. Fetus turned for delivery. Woman died of puerperal inflammation.
9	1778	Dr. J. R. Sigault	Paris	3 in.	Recovered	Lost	Wife of a soldier; had had 7 children, all of whom died; one by embryotomy; pubes ossified, and was saved in labour. Fetus turned for delivery. Woman died of puerperal inflammation.
10	1778	Dr. J. R. Sigault	Paris		Recovered	Lost	Wife of a soldier; had had 7 children, all of whom died; one by embryotomy; pubes ossified, and was saved in labour. Fetus turned for delivery. Woman died of puerperal inflammation.
11	Nov. 13, 1778	Dr. J. R. Sigault	Paris		Recovered	Lost	Wife of a soldier; had had 7 children, all of whom died; one by embryotomy; pubes ossified, and was saved in labour. Fetus turned for delivery. Woman died of puerperal inflammation.
12	Nov. 13, 1778	Dr. J. R. Sigault	Paris	23 or 24 l.	Died	Lost	Wife of a soldier; had had 7 children, all of whom died; one by embryotomy; pubes ossified, and was saved in labour. Fetus turned for delivery. Woman died of puerperal inflammation.
13	Feb. 12, 1779	Dr. Duret	Brest, France		Recovered	Lost	Wife of a soldier; had had 7 children, all of whom died; one by embryotomy; pubes ossified, and was saved in labour. Fetus turned for delivery. Woman died of puerperal inflammation.
14	June 30, 1779	Dr. Van Damme	Racqueghem, Fr.		Recovered	Alive; recovered	Wife of a soldier; had had 7 children, all of whom died; one by embryotomy; pubes ossified, and was saved in labour. Fetus turned for delivery. Woman died of puerperal inflammation.
15	July 18, 1779	Dr. Alphonse Le Roy	Paris	2 in. 8 l.	Recovered	Alive	Wife of a soldier; had had 7 children, all of whom died; one by embryotomy; pubes ossified, and was saved in labour. Fetus turned for delivery. Woman died of puerperal inflammation.
16	July 21, 1779	Dr. Alphonse Le Roy	Paris	2 in. 0 l.	Recovered	Alive	Wife of a soldier; had had 7 children, all of whom died; one by embryotomy; pubes ossified, and was saved in labour. Fetus turned for delivery. Woman died of puerperal inflammation.

Symphysiotomy Table 1, continued.

No.	Date.	Authority.	Locality.	Conjugate diameter.	Result to woman.	Result to child.	Remarks.
17	Jan. 1780	Dr. Du Cambon	Mons, Belgium		Recovered	Alive	A second operation performed upon the subject of case 4. Woman 42, a primipara, and in labour 3 days. Operation by a herniotomy under the direction of Prof. Francesco Civallotti, who had been previously employed in the same manner, and without any untoward result. The child was delivered in a head-down position, and the placenta was delivered without any untoward result. Woman recovered in 35 days.
18	Aug. 9, 1780	Directed by Prof. F. Civallotti, of the University of Padua	Ugento, Apulia, Italy		Recovered	Alive	
19	Dec. 5, 1781	Dr. Du Chaulfuy	Lyon	1 in. 7 l.	Died	Lost	Inclination unaltered; knife cut off the end of right pubic bone. Europe used, then turning. Woman died in 32 hours, from fatal phlogia.
20	Dec. 24, 1781	Dr. Antonio Lavagnolo	Genoa	2 in. 5 l.	Died	Alive; soon died	Female child, thought dead, was extracted with crochets, and lived several hours. Woman died in 17 days, external and internal parts gangrenous.
21	Sept. 4, 1782	Mr. John Wetchman	Kington, England	2 1/2 in.	Died	Lost	Case of malaccetion; woman reduced from 5 ft. 6 in. to 4 ft. 4 in. labour; pulse 110; fetus putrid; long labour. First and second operations by symphysiotomy in Great Britain.
22	Mar. 1783	Dr. Giovanetti	Borgo d'Arza, Italy		Recovered	Lost	Partial symphysiotomy; woman recovered; in labour 3 days; pulse but slightly elevated; no phlogia.
23	Oct. 20, 1783	Dr. Johannes G. Damon	The Hague		Recovered	Lost	Was once before delivered of a putrid child; phlogia appeared by a razor and a common knife; head escaped as the patient was expiring.
24	Nov. 1783	Dr. V. (name withheld)	Paris		Died	Lost	Third labour; delivered by turning; patient recovered by 17th day.
25	Aug. 7, 1784	Dr. Alphonse Le Roy	Paris	3 in.	Recovered	Alive; soon died	Woman had borne 4 children before; the first died at 15 months; second was a foetus, and lost; third delivered by the crochets. Pubic bone was cut through in the pubiotomy, as in case 21; fetus delivered by the feet. Biparietal diameter 3 1/2 inches.
26	Mar. 12, 1785	Dr. Alphonse Le Roy	Paris	Over 3 in.	Recovered	Alive; soon died	Cardinal's case; woman delivered by her pelvis. Cardiac pulse; in labour 3 days; no phlogia. Woman died on 15 and 27. Biparietal diameter 3 in. 5 lines. Woman died on 15th day; intra-pelvic parts found gangrenous.
27	Apr. 17, 1785	Dr. Demathis	Paris	2 in. 6 l.	Died	Lost	A second operation performed up in the subject of case 21. Woman recovered, but died on 15th day from phlogia.
28	Apr. 21, 1785	Dr. Alphonse Le Roy	Paris	2 in. 6 l.	Died	Alive	Woman 28; in labour 30 hours; child lived less than half an hour. Woman delivered in 40 days; had a urinary fistula in lower part of clitoris.
29	Aug. 11, 1785	Dr. J. C. Dymen	The Hague		Recovered	Alive	Symphysis failed to unite. Woman still living, and able to walk twenty years later; was employed as a messenger.
30		Dr. De Cambron	Mons, Belgium		Recovered	Alive	Case of malaccetion; woman died of anasarca in 40 days; symphysis united in 30 days.
31		Dr. Du Cambon	Mons, Belgium		Recovered	Alive; soon died	
32	Jan. 21, 1786	Dr. Verdier Ducloux	La Ferté Bernard, France	2 in. 7 l.	Died	Lost	
33		Dr. Du Cambon	Mons, Belgium		Recovered	Alive; soon died	
34	Rep. in 1801	Prof. Domenico Ferraro	Naples		Recovered	Lost	
35	1807	Dr. Johannes Van Munster	Symingen, Holland		Died	Lost	
		Brano Amantea	Naples		Died	Lost	

1 Caseus credit. Imbert of Lyons, 1833, with originating this method of operating, done 52 years before.

Symphysiotomy Table I, concluded.

N ^o .	Date.	Authority.	Locality.	Conjugate diameter.	Result to woman.	Result to child.	Remarks.
36		Prof. Dubois	Paris		Recovered	Lost	Patient had been once before delivered.
37	1853	Dr. Vermandels	Bourg, France		Recovered	Lost	Woman 30; second labour. Craniotomy and turning tried before opening the pelvis; then turning and forceps.
38	Oct. 24, 1869	Prof. Dubois and Gardien	Paris		Recovered	Alive	Woman 27; in her third labour.
39	1811	Dr. Paolo Asallini	Paris		Recovered	Alive	W. P. A. is credited with two cases, both saved.
40	1811	Dr. Vermandels	Bourg, France		Recovered	Lost	Woman 27; in her third labour. Craniotomy and turning tried before opening the pelvis; then turning and then forceps used after the pelvis was opened.
41	1815	Prof. Gennaro Galbati	Naples	2 in. 4 l.	Died	Lost	Woman died of putrid fever in 15 days; child lived 43 hours; foot presentation.
42	1817	Dr. Giacomo Giannini	Milan		Recovered	Alive	Woman died of putrid fever in 15 days; child lived 43 hours; foot presentation.
43	Jan. 13, 1816	Dr. G. Giannini	Milan	2 in. 3 l.	Died	Lost	Woman died of putrid fever in 15 days; child lived 43 hours; foot presentation.
44	1817	Dr. G. Giannini	Milan		Died	Lost	Woman died of putrid fever in 15 days; child lived 43 hours; foot presentation.
45	1817	Dr. G. Giannini	Belligny, France		Recovered	Alive	Woman died of putrid fever in 15 days; child lived 43 hours; foot presentation.
46	1	Dr. Rollay	Pinopol, France	3 in.	Died	Lost	Child turned for delivery. Patient died of syncope in an hour and a half.
47	?	Dr. Brothling, Jr.	?		Recovered	Lost	Woman had previously borne 2 children. Before incising the symphysis, the arm had been amputated and fetus destroyed.
48	1818	Dr. Prillavitz	Holland		Recovered	Lost	Woman recovered in 22 days; her health was good afterwards.
49	1818	Prof. G. Galbati	Naples	2½ in.	Recovered	Alive	Woman recovered in a month; her pelvis was deformed by <i>valvulation</i> .
50	1820	Prof. G. Galbati	Naples		Recovered	Lost	Woman recovered in a month; her pelvis was deformed by <i>valvulation</i> .
51	1821	Prof. Mandel	Naples	2½ in.	Recovered	Alive	Patient was well in 40 days.
52	1820	Prof. Mandel and Mandel	Naples	2½ in.	Recovered	Lost	Bleech presentation; fetus moribund when delivered; puerperal consolidation in 35 days.
53	1828	Prof. G. Galbati	Naples	2½ in.	Recovered	Alive	Woman died in 5 days of gangrene of the vagina, etc.
54	1825	Dr. J. Pasquale Catellana	Naples	2½ in.	Died	Lost	Patient recovered in 35 days. A small piece of bone exfoliated from the pelvis.
55	1831	Dr. Francesco Petrucci	Naples	2½ in.	Recovered	Alive	Fetus died the first day from injury by forceps. Woman affected with leucorrhoea of urine.
56	1835	Prof. G. Galbati	Naples	2½ in.	Recovered	Lost	Fetus delivered by the forceps.
57	1835	Prof. G. Galbati	Naples		Recovered	Lost	(Twins.)
58	1833	Prof. G. Galbati	Naples		Died	Lost	
59	1830	Prof. G. Galbati	Naples	2½ in.	Died	Lost	
60	1830	Prof. G. Galbati	Naples	2½ in.	Recovered	Alive	
61	1830	Prof. G. Galbati	Naples		Recovered	Alive	
62	1837	Prof. G. Galbati	Naples		Died	Lost	
63	1833	Prof. G. Galbati	Naples		Recovered	Alive	
64	1839	Prof. G. Galbati	Naples		Died	Lost	
65	1840	Prof. G. Galbati	Naples		Died	Lost	
66	1840	Prof. G. Galbati	Naples		Died	Lost	
67	1866	Dr. Caccavelli	Naples	3 in.	Recovered	Alive	
68	1833	Prof. Enrico Jacolucci	Naples	2½ in.	Recovered	Lost	Inferior retract contracted; fetus extracted with forceps; woman recovered in 40 days.
69	1836	Prof. E. Jacolucci	Naples	2½ in.	Recovered	Alive	No accident, well in a month.
70	1858	Prof. E. Jacolucci	Naples	2½ in.	Died	Lost	Woman died in 12 days of pelvic cellulitis; was 8 months pregnant.

My thanks are due in connection with the preparation of this historical table to Prof. Alfonso Corradi, of Bologna, who kindly sent me his voluminous quarto history of obstetrics in Italy, completed in 1875; and to Dr. J. Stockton Hough, formerly of this city, for the privilege of examining his collection of rare symphysiotomy pamphlets. Those who make researches in old obstetrical matters often bless the memory of the late Prof. Charles D. Meigs, who, when in Europe some thirty years ago, made a collection of old works pertaining to his special branch, which are now accessible to the student, in the libraries of the Pennsylvania Hospital and College of Physicians, of Philadelphia.

Summary of the 50 Recent Neapolitan Cases.—Women saved, 40; died, 10. Children born alive, 41; dead, 9. Vertex presentations, 45; shoulder, 2; breech, 3. Conjugate diameters, $2\frac{3}{8}$ inches in 2 cases; $2\frac{5}{8}$ inches in 15; $2\frac{3}{4}$ inches in 5; $2\frac{7}{8}$ inches in 16; and $3\frac{1}{8}$ inches in 12 = 50. The same in the fatal cases, $2\frac{3}{8}$ inches in 5 (lost out of 15); $2\frac{7}{8}$ inches in 4 (lost out of 16); and $3\frac{1}{8}$ inches in 1 (lost out of 12) = 10. The fetuses perished in all of the shoulder (2) and breech presentations (3), and in 4 of the 45 vertex presentations. Nine of the 10 women lost were delivered of living children; and 8 of the 9, who bore dead children, recovered. In only one instance did both mother and child perish, and this was a case of breech presentation in one of the largest ($3\frac{1}{8}$ conj.) pelves. The woman was operated upon on the fourth day of labour. Cases 8-19 and 24-28 were each operated upon twice successfully, making in all six double operations, or six subjects twice operated upon, on record.

The two tabular records contain, as stated, 120 cases. To these may be added a second successful operation by Assalini, and one each for Bal-lentani and Marescotti, all of Lombardi; the last two prior to 1812; one successful, for Ettore Piccinnini, of Asti, 1871; two for Prof. Novi, of Naples, to be hereafter referred to more particularly; and one to Prof. Morisani, of Naples, 1880. These seven additional cases increase the list to 127, with 92 recoveries. Besides these, are operations, one each, said to have been performed in Constantinople, Bannières-en-Artois, France, and Frankfort-on-the-Main, in the last century, or very early in this, results not given; and no doubt others entirely lost to history.

Italy alone has a credit of 87 symphysiotomies, with 65 women saved, against perhaps as many as 50 for the rest of Europe, more than half of which were in France. The first operation performed in Italy has generally been credited to Dr. Antonio Lavaguigno, of Genoa (No. 20 of table 1), in 1781; and the first in the *Ospedale dei Incurabili*, to Prof. Domenico Ferrara (No. 33 of table 1), in 1787; but Prof. Ottavio Morisani, of the same hospital, has just made¹ claim, in a foot-note to a translation of Cazeaux and Tarnier's Midwifery, that Ferrara performed the

¹ October, 1882.

2. A Table of 50 recent Neapolitan Symphysiotomies, showing a greatly diminished mortality.

No.	Date of operation.	Hospital.	Name of patient.	Age	Conj. diam. in. millie.	Proseca-tion.	Result to woman.	Result to child.	Cause of death of woman.	Remarks.
1	Jan. 5, 1869	For incurables	Anna Esposito	29	3 1/2	Vertex	Recovered	Alive		
2	Nov. 17, "	"	Anna Esposito	39	3 1/2	Vertex	Recovered	Alive		
3	Jan. 23, 1867	"	Bridgetta Pugliese	40	3 1/2	Breech	Died	Dead	Metropertionitis	Admitted on the fourth day of labour.
4	June 29, "	"	Giustina Altanuello	38	3 1/2	Vertex	Recovered	Alive		Admitted when in labour.
5	Dec. 29, "	"	Maria Angiano	29	2 1/2	Vertex	Recovered	Alive		
6	Oct. 20, "	"	Ferdinando Esposito	29	3 1/2	Breech	Recovered	Dead	Metropertionitis	Membranes ruptured 14 hours.
7	Jan. 28, 1868	"	Isabella Bruno	23	3 1/2	Vertex	Recovered	Alive		Had been 45 hours in labour when admitted.
8	Aug. 4, "	"	Vincenza Romano	23	2 1/2	Vertex	"	"		
9	Aug. 4, "	"	Anna Bongiovanni	21	2 1/2	"	Died	"	Metropertionitis	
10	Oct. 4, "	"	Nunzia Nucera	31	2 1/2	"	Recovered	"		
11	Dec. 14, "	"	Terese Guasto	31	2 1/2	Shoulder	Recovered	Dead		
12	Dec. 14, "	"	Concetta Esposito	31	3 1/2	Vertex	"	Alive		
13	Feb. 6, 1869	"	Concetta Esposito	17	2 1/2	"	"	"		
14	June 1, "	"	Angela Viola	23	2 1/2	"	Died	"	Metropertionitis	
15	June 6, "	"	Flora Maria Tarallo	21	2 1/2	"	Recovered	"	Peritonitis	
16	July 5, "	"	Concetta Borbonico	21	2 1/2	"	"	"		
17	July 6, "	"	Maria Cantolino	21	2 1/2	"	"	"		
18	Aug. 2, "	"	Concetta Esposito	33	3 1/2	"	"	"		
19	Apr. 27, 1870	"	Vincenza Conci	33	3 1/2	"	"	"		Second operation; see Case 8.
20	Oct. 7, 1871	"	Rosa Barghese	22	2 1/2	"	"	"		
21	"	"	Rosa Fauro	22	2 1/2	"	"	"		
22	"	"	Rosa Monte	22	2 1/2	"	"	"		
23	"	"	Terese Arnesa	22	2 1/2	"	"	"		
24	"	"	Maria Battagliari	22	2 1/2	Shoulder	"	Dead		A vesico-vaginal fistula produced; cured by an operation.
25	"	"	Maria Battagliari	22	2 1/2	Vertex	"	Alive		
26	Feb. 7, 1873	"	Maria Battagliari	22	2 1/2	"	"	"		
27	Aug. 30, "	"	Terese Arnesa	22	2 1/2	"	"	"		There were 29 operations and 6 deaths, up to Sept. 25, 1872 (Goratti).
28	Mar. 11, 1874	"	Rosa Alumba	22	2 1/2	"	"	Dead		Second operation; see case 21.
29	Feb. 20, 1875	"	Maria Lecce	22	2 1/2	"	"	Dead		
30	Feb. 20, 1877	"	Giuseppina Crispo	20	2 1/2	"	"	Alive		
31	May 13, "	"	Maria Anselmo	20	2 1/2	"	"	"		
32	May 13, "	"	Giuseppina Crispo	20	2 1/2	"	"	"		
33	Feb. 28, "	"	Carmina De Luca	27	2 1/2	"	"	"		In labour 30 hours when admitted.
34	Aug. 7, "	"	Vincenza N.	27	2 1/2	"	"	"		
35	Aug. 7, "	"	Maria Finello	27	2 1/2	"	"	"		
36	Aug. 7, "	"	Giuseppina Crispo	27	2 1/2	"	"	"		
37	Nov. 1, "	"	Isabella Esposito	29	2 1/2	"	Died	"	Peritonitis	
38	Apr. 27, 1878	"	Isabella Esposito	29	2 1/2	"	Recovered	"	Eudocarditis.	Lived 32 days after the operation.
39	Dec. 10, "	"	Flora Maria Carrano	29	2 1/2	"	"	"		
40	Dec. 10, "	"	Vincenza Nunziato	22	2 1/2	"	"	"		
41	Feb. 6, 1879	"	Maria Ippolito	23	2 1/2	"	"	"		
42	Mar. 8, "	"	Nunzia Isello	22	2 1/2	"	"	"		
43	Oct. 20, "	"	Concetta Esposito	22	2 1/2	"	"	"		
44	Oct. 20, "	"	Rosa Bocca	22	2 1/2	"	"	"		
45	Dec. 9, "	"	Giuseppina Petralicchi	23	2 1/2	"	Died	"	Diphtheritic endometritis and Septic metropertionitis.	The disease prevailed at the time in the hospital. She had been in labour 20 hours when admitted.
46	Jan. 6, 1880	For incurables	Angela Esposito	20	2 1/2	"	Recovered	Alive		
47	Jan. 10, "	"	Cristina Pastore	22	2 1/2	"	"	"		
48	Oct. 20, "	"	Rosa Rosati	22	2 1/2	"	"	"		
49	Dec. 21, "	"	Carolina Passalacqua	22	2 1/2	"	"	"		
50	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	Died	"		
51	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
52	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
53	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
54	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
55	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
56	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
57	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
58	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
59	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
60	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
61	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
62	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
63	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
64	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
65	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
66	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
67	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
68	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
69	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
70	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
71	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
72	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
73	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
74	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
75	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
76	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
77	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
78	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
79	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
80	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
81	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
82	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
83	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
84	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
85	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
86	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
87	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
88	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
89	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
90	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
91	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
92	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
93	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
94	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
95	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
96	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
97	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
98	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
99	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
100	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
101	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
102	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
103	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
104	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
105	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
106	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
107	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
108	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
109	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
110	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
111	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
112	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
113	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
114	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
115	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
116	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
117	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
118	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
119	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
120	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
121	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
122	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		
123	Dec. 21, "	Obstetric Clinic		32	2 1/2	"	"	"		

operation in that institution in 1774, which would have been three years earlier than that of Sigault. This statement is based on that of Prof. Cattolica, which was set aside by Prof. Corradi, in his History of Obstetrics in Italy. Galeotti published an obstetrical treatise in Naples, in 1787, in which he refers to the operation of Ferrara of that year, as the second in Italy, and that of Lavaguigno as the first. This is, in one sense, also an error, as the case of Giovanetti, 1783 (No. 22, table 1), is between the two. Ferrara is said by Novi, of the same hospital, not to have been made a professor until 1777, three years after he is claimed to have operated. Had Galeotti been in error, he would doubtless have been corrected by Ferrara as to the time of his first operation.

According to Penchienati and Brugnone, 1866, there were 34 symphysiotomies in Europe from 1777 to 1785, with 23 mothers and 11 children saved. Baudelocque gave 25 as the number of operations in the first five years. Churhill, in 1841, gave the record of cases as 49, and deaths as 16. It will now be seen by my record that at that time there had been at least 69, with 25 deaths. After the operation was inaugurated in the Hospital for Incurables in Naples, ninety-five years ago, it was occasionally resorted to at longer or shorter intervals, alternating with the Cæsarean section as the cases appeared to require it, until finally there were but 9 symphysiotomies in a period of thirty years (1836-66), saving only 4 women and 3 children. During the eight years prior to 1866, in which the operation was revived, there was not a case. After the resuscitation, there was such a change of success, that the operation advanced in the estimation of the hospital staff, and has, since January, 1866, been performed more than fifty times.

My second table begins with the resuscitation in Naples, and up to the present time, as far as I have been able to ascertain, there have been 53 operations in that city, saving 43 women and 42 children. In the Hospital for Incurables, where nearly all of these women were operated upon, they have had a bitter experience with the Cæsarean section, losing 25 out of 27 women between 1791 and 1875, although all of the children were removed alive. With an abundant rachitic population requiring relief in parturition, is it to be wondered at that the obstetric surgeons of Naples should endeavour to revive and improve the method of Sigault, notwithstanding its general condemnation, and their own very imperfect success? When we read the record of success, in table No. 2, we are inclined to ask, can this be the "unscientific" and "unjustifiable" operation of Sigault, and if so, what has made the great change in the mortality of the women and children, particularly of the latter? And these are not the only improvements claimed, as will be shown by an examination of the causes of death in the two tables, and attention to the following answers to an interrogatory letter, returned by Professor O. Morisani, of the University of Naples.

1. All of the fifty operations (in table 2) were performed upon rachitic subjects, whose pelves were generally flattened antero-posteriorly. In four or five instances the pelves were simply dwarfed in dimensions. There was no case of rostrate pelvis, as *malacosteon* is very rarely met with in Naples.

2. Version was not resorted to except in the transverse positions. The forceps were applied in about one-fourth of the cases.

3. The separation at the pubes amounted to about 2 inches (50 mm.), which was obtained without any effort, and without producing any lesion of the sacro-iliac synchondroses.

4. The immovable dressing secured the firm union of the symphysis pubis in all the cases that recovered.

5. The women had good health after the operation.

6. There were no malformed infants. Nearly all of the children were sent to the Foundling Hospital to be taken care of.

7. *Phlegmasia alba dolens* did not occur in any of the women.

8. There were no pelvic lesions left, as a sequel of the operation.

9. Vesico-vaginal fistula occurred in but one case, and this was easily cured by an operation.

The answer to No. 8, I presume, is an oversight, as table 2 records a death from "*iliac phlegmon*," case 5.

Table No. 2 was prepared for me in Naples after a form sent; but by an error in transcribing, the heading of the column arranged for the operators (*operatori*) was changed to *operata*, and the names of the patients made to take those of the operators whom I designed to credit. These, however, were mainly Professors Morisani and Novi, already mentioned.

The operation in Naples has been performed with great carefulness. The section is made subcutaneously with the probe-pointed and sickle-shaped (*falcetta*) bistoury of Galbiati. An incision is made above the pubes, somewhat after the method of Ferdinando Carbonai (1841), of Florence, and the knife of Galbiati slowly passed behind the symphysis until it reaches the pubic arch, when its cutting edge is brought to bear upon the ligaments, and the parts are divided from below upward. The pelvis is not forced open, neither is the fetus turned or dragged upon, but where the head presents, the case is left mainly to nature. In about one case out of four the forceps are applied as an assistant. The incised part is treated antiseptically, and by irrigation if in warm weather; and as soon as convenient the ossa pubes are kept in apposition by an immovable apparatus, to secure an early union of the severed parts.

To avoid the much-dreaded Cesarean operation, Professor Enrico Jacolucci, of Naples, in 1867, proposed to combine the induction of premature labour with symphysiotomy in one class of extremely deformed cases, and with craniotomy and cephalotripsy in another and still more deformed class. Acting upon his suggestions, Professor Novi has performed

one operation of each class, and Professor Morisani one of the former, as follows:—

1. The first operation was performed by Prof. Novi, in the Hospital for Incurables, upon Louisa Attiola, having a conjugate of $2\frac{1}{2}$ inches (54 mm.). Labour was induced in the seventh month, the fœtus presented in the second position of the right shoulder, was turned and delivered, but lived only an hour. The woman recovered in 50 days.

2. The second operation by him was performed at term, upon Rosa Meglio, on September 8th, 1872. This woman had a conjugate of less than 2 inches (49 mm.), and the fœtus was dead. After opening the symphysis, the head was perforated and then crushed, after which it was delivered with the body of the fœtus; the woman recovered in 42 days. In estimating the propriety of this mode of operating, it must be borne in mind that one of the two successful gastro-hysterotomies in Naples out of 27 was performed by Prof. Novi, in 1865, and that in 1871, his only former case had died of peritonitis in three days. He doubted the propriety of risking the Cæsarean section for the delivery of a dead child.

3. The third operation is one of much interest, and was performed by Prof. O. Morisani, upon Lucia Esposito, a rachitic dwarf, 20 years old, and 3 ft. $7\frac{3}{4}$ inches high, having a conjugate of $2\frac{3}{4}$ inches ($5\frac{1}{2}$ cm.). Of this woman I have in my possession two full-length photographs, and have received from Dr. Rafaele, in pamphlet, the following description: "Head large; lower jaw elongated; teeth lost or decayed; right shoulder-blade and ribs beneath more salient than the corresponding parts on the left side; left lateral inclination of the thorax, with a convexity in the right dorsal portion; compensative scoliosis of the lumbar portion; back very hollow; and right natis much more prominent than the left." This woman entered the *Clinica Ostetricia* on May 15th, 1880, in the seventh month of her pregnancy, and when the extent of her pelvic deformity was ascertained, it was determined to bring on labour in the first week of the eighth month. This was accordingly done on June 9th, and by the 11th labour had sufficiently advanced to permit of the performance of symphysiotomy, which was executed, after the manner already described, by Prof. Morisani assisted by Prof. Novi and others. When the symphysis was divided, the fœtal head which presented by the vertex in the first position began to descend, passed into the cavity of the pelvis, and after some delay at the perineum was extruded from the vulva. The fœtus soon began to breathe regularly; it was $15\frac{3}{4}$ inches long, and weighed $4\frac{1}{4}$ pounds; occipito-frontal diameter $3\frac{1}{2}$ inches; occipito-mental $4\frac{3}{4}$; biparietal 3 inches (7.5 cm.); and bi-temporal $2\frac{3}{4}$ inches. The child when three days old was sent to the *Casa della Annunziata*, a foundling hospital.

The wound was dressed with a compress, kept moist by a drainage tape leading from a vessel of hæmostatic water, and by the end of a week the pelvis was secured by an immovable apparatus, an opening being left

over the wound. In 34 days union was complete, and in forty days the patient left her bed, being ready soon afterward to be presented at the clinic, having no pain or inconvenience in walking.

Symphysiotomy would appear to present, according to its history, two eras having in each very opposite characters, as shown by their relative mortality in women and children, and the effects on the health of the surviving subjects. Thanks to the opposition of Baudelocque and many obstetrical writers of the close of the last and beginning of the present century, I have been enabled to procure and present the results of the early operations from 1777 to 1815, by which it will be seen that not only was the mode of delivery very fatal to the women, and still more so to the children, but the opening of the pubes was made to produce such injuries to the sacro-iliac connections as to set up an inflammation ending in gangrene and death. It is clearly demonstrated by Baudelocque that the early operators were often incapable of measuring the interior of the pelvis with any accuracy, and that, in many instances, women capable of bearing unmutilated and in some cases living children were subjected to the operation. By comparing the causes of death in tables 1 and 2, it will also be noticed that they are altogether different. In the first we have generally death from pelvic injuries, and in the second from peritonitis, metro-peritonitis, and other forms of inflammation, such as follow cases of labour, not always instrumental or traumatic. Italian hospital obstetricians, in the regions where pelvic deformities are most numerous, have become of necessity skilled in the use of pelvimeters of various forms, and, from constant practice, are enabled to make very reliable measurements in parturient subjects. They have also been forced to adopt every expedient of modern surgery in after-treatment to diminish the mortality in their old maternities, which were formerly little better than pest-houses in the proportion of deaths. The effect of antiseptic measures and greater cleanliness has been shown by the results of the Porro and other capital operations in large lying-in institutions, and in nothing more than the entire change of results in the two maternities of Naples in which the old and new pubic sections have been performed. Perhaps no tabular record of Italian surgery is so much to the point in exhibiting the possible reversion of results from unfavourable to favourable as that prepared upon the first, second, and third hundred ovariectomies, by Dr. Peruzzi, of Lugo, and reviewed in abstract by me in this Journal. From having lost nine of the first operations in succession before one success, they have gradually improved in results until now the mortality is reduced to a moderate percentage, and the end of reduction has evidently not yet been reached. It is not strange, then, that they have been able to improve materially in their results in symphysiotomy in the old Ospedale dei Incurabili of Naples, where fifty years ago, and more recently, Galbiati, whose knife they now use, was so unsuccessful in

his results, as shown towards the close of table 1. Had this operation been all that William Hunter, Baudelocque, Churchill, and many others believed it to be, scientifically considered, it would have been beyond the possibilities for Morisani and Novi to have demonstrated its feasibility upon the living subject as they have done. There is certainly more in the theory of pelvic mobility in pregnant women than was demonstrated by Hunter upon the dead subject, the much more extensive test upon the living having shown the fallacy of his. It can easily be shown how little is gained in the sacro-pubic diameter for each inch of pubic separation; but this does not appear to present so great an obstacle in practice as might be supposed. The pelvis gains in its transverse and oblique diameters, and the head, when not hurriedly forced, moulds itself to the space obtained and slowly passes through. At all events, the fact remains that 42 out of 46 fetuses presenting by the vertex were in some manner delivered alive, in Naples, since January, 1866, under pubic section, with a saving of 80 per cent. of the mothers. Other hospitals, and obstetricians in other countries in private practice, may not be able to accomplish as much, but the fact of the possibility remains. The Porro operation in two hospitals in Vienna and Milan has been managed with such care and skill that 14 out of 19 women have been saved (73 per cent.), and all of the children; but the rest of the world out of 77 cases has only saved 30, or 38 $\frac{7}{8}$ per cent., with 55 children. Two of these seventy-seven operations were performed in Naples in 1881 and 1882, and both women, with one child, were lost. The Porro operation thus far has saved 44 out of 96 women, and 74 children, or 45 $\frac{5}{8}$ per cent. of the mothers, and 77 $\frac{1}{2}$ per cent. of the children. Prof. Morisani, before the International Medical Congress of 1881, was inclined to compare the symphysiotomies of the Neapolitan hospitals with the *general* work under the Porro operation, which we think unfair to the latter. Let him measure their 50 cases with those of Santa Caterina of Milan and the Krankenhaus of Vienna, and he will show a saving of 80 per cent. of women and 82 per cent. of children, against 73 per cent. of women and 100 per cent. of children.

But in one sense the two operations are not to be compared. Symphysiotomy can never, with every advantage of care, skill, and climate, except in a limited degree, be made to take the place of the Cesarean and Porro-Cesarean operations, as its advocates are not inclined to recommend it in cases having a conjugate of less measure than 67 millimeters, or 2 $\frac{5}{8}$ inches. Conscious of this fact, some of the early advocates of symphysiotomy endeavoured to modify the pubic section, so as to make it possible to deliver in cases of extreme deformity, and to this end devised the operation known as *bi-pubiotomy*, which in its fatal results did much to hasten the downfall of the more simple method. The operation originated with Prof. Galbiati of Naples, who performed it on a dwarf 3 $\frac{1}{2}$ feet high, on March 30th, 1832. As this woman had a conjugate of

only an inch, he devised the plan of opening her pelvis, by a subcutaneous section with an Aitkin chain-saw, cutting the horizontal and descending rami of the ossa pubes on either side, so as to be able to open the collapsed superior strait, by lifting the anterior wall of the pelvis from the posterior, as we open a bellows. He succeeded in delivering the woman of a dead fœtus, but such was the injury produced by stretching the tissues over the severed bones, that she died in agony on April 3d, four days later, her vagina, vulva, and surrounding tissues being all gangrenous. Not contented with this disastrous test, Dr. Nanzianti Ippolito tried the same plan in the Hospital for Incurables in the winter of 1842-3 with a similar fatality. Thus ended the attempt to make a pelvic section do the work of a gastro-hysterotomy in cases of excessive pelvic deformity.

Symphysiotomy has found a home in Italy for two reasons, viz., the number of parturient women demanding relief on account of pelvic deformity, and the opposition of the Papal church to the destruction of unborn infants. In a country like Great Britain or the United States, cases with the conjugate limits in table No. 2, would be delivered by craniotomy or cephalotripsy, as the life of the fœtus is considered of no value where that of the mother is in danger. But although the general teaching of our text books is to this effect, there are exceptional cases in which there should be an effort made to save the fœtus. If symphysiotomy can be performed with safety to the mother, why should it not be employed in cases where the pelvis is just a little too small to admit of the fœtus being born alive, and where fœtus after fœtus has been sacrificed to save the mother? Such cases could be relieved and saved by the induction of premature labour, but the opportunity to do this at the proper time is generally wanting, and patients are often unwilling to submit to it. I have seen a woman lose child after child, simply because her pelvis was small, and the foetal heads large; and yet occasionally deliver herself where the head was small enough to pass. There have been as many as sixteen children sacrificed in the successive labours of one lady in this city, each head having been locked in her pelvis, which was small but not deformed: a seventeenth was saved, by an accident inducing labour, when she was 8 months pregnant. Now is there not a field for symphysiotomy in such cases, if the operation can be performed with the small amount of risk claimed in table 2? Has the fœtus no claim upon us for its life, and must we always destroy it in the interest of the mother? I believe with the late Dr. Thomas Radford of Manchester, Eng., that the fœtus has more title to life than we generally accord it.

Symphysiotomy is by no means the "very simple" operation that writers have claimed it to be. As deformed pelves are frequently asymmetrical, it is difficult to define the exact centre of the symphysis pubis, just as it is to determine in an ovariectomy the position of the *linea alba*. If the end of one os pubis is sliced off as has several times been done, the piece

excised will become carious and keep up a discharging fistula until it is expelled or removed. An error in striking the line of the cartilage will also delay very materially the process of union. The operation is best adapted to cases of rachitic deformity, or to symmetrical dwarfing of the pelvis, and should be avoided where there is any reason to believe that the deforming disease may have produced sacro-iliac ankylosis on one or both sides. It would therefore be improper to operate in a case of Naegelé's oblique pelvis, or Robert's pelvis; or of the oblique pelvis produced by coxalgic ankylosis of one side, in which the os innominatum is undeveloped, flattened, and apt to be united by bone to the sacrum.

The result of the operation will often be determined by the relative size of the foetal head; and the impossibility of ascertaining its dimensions before delivery, constitutes another of the difficulties in the way of operating. A dwarf may carry in utero a fœtus which is out of all proportion to her own size (see case 12, table 1), and the possibility of this should be suspected, if the father is of large proportions. Rachitic dwarfs are usually illegitimately impregnated, and although they may attempt to conceal this fact, the size of the father of the child can generally be ascertained, at least so far as to his being a small or tall man. A woman of 4 feet 4 inches high, bore in this country a fœtus of 14 pounds weight; and one of 3 feet 2 inches high, a child weighing 9 pounds: both were removed by the Cæsarean section. It is evident that in such cases, the operation of Sigault must be a failure. In the lower animals, the impregnation of a small female by a much larger male is not infrequently fatal to her in labour from the excessive size of the fœtuses. A few years ago a leopardess died in labour in the garden of the Philadelphia Zoological Society from this cause. She had been brought up in captivity and her skeleton was of small size, while the male was a large trapped animal caught after maturity; three cubs were extruded, and four larger ones remained in utero, where they were found *post mortem*.

Where premature delivery at the earliest viable period is impossible, it may be advisable to bring on labour and then open the symphysis as in the Morisani case already related. The fatality of such cases under craniotomy and the Cæsarean section influenced the Neapolitan obstetricians to offer this method as one less dangerous to the mother, as well as saving the fœtus: the gain of a fraction of an inch may decide the question between its life and death.

Version by the feet was the common practice of the early symphysiotomists, and no doubt the cause of serious injury to the mother and the death of the fœtus in many cases. The abandonment of this practice has very materially reduced the fatality of the operation to the fœtus, and the amount of strain put upon the sacro-iliac synchondroses. Traction and version by the feet are confined, in Naples, to non-cephalic presentations, as of the shoulder, body, breech, and feet.